

SCAR markers linked with disease resistance traits in common bean (*Phaseolus vulgaris*)

Updated: October, 2007

SCAR Name	Marker of Origin	Pathogen	Size (bp) / orientation	Sequences of SCARS	Tagged Locus	LG	Reference
SAP6	AP6	Common Bacterial Blight (CBB)	820 cis	F - GTC ACG TCT CCT TAA TAG TA R - GTC ACG TCT CAA TAG GCA AA	Major QTL (GN#1 sel 27)	B10	Miklas et al., 2000b,c
BAC6	BC409	CBB	1250 cis	TAG GCG GCG GCG CAC GTT TTG TAG GCG GCG GAA GTG GCG GTG	Major QTL (GN#1 sel 27)	B10	Jung et al., 1999
SU91	U9	CBB	700 cis	CCA CAT CGG TTA ACA TGA GT CCA CAT CGG TGT CAA CGT GA	Major QTL (XAN 159)	B8	Pedraza et al., 1997
LG5 syn. BC420	BC420	CBB	900 cis	GCA GGG TTC GAA GAC ACA CTG G GCA GGG TTC GCC CAA TAA CG	Major QTL (XAN 159)	B6	Yu et al., 2000
R7313		CBB	700 cis	ATT GTT ATC GTC GAC ACG AAT ATT TCT GAT CAC ACG AG	Major QTL (OAC 88-1)	B8	Bai et al., 1997 Beattie et al., 1998
R4865		CBB	950 cis	TCC AAA GCC ATT CTA GTT CAG CTA CTT TCA AAC TGG G	Major QTL (OAC 88-1)		Bai et al., 1997 Beattie et al., 1998
SB10	B10	Halo bacterial blight (HBB)	525 cis	CTG CTG GGA CAA TCA CCA AGT C CTG CTG GGA CTC TCT TAC	<i>Pse-1</i>	B4	Fourie et al., 2004

SW13	W13	Bean Common Mosaic Virus (BCMV) & HBB	690 cis	CAC AGC GAC ATT AAT TTT CCT TTC CAC AGC GAC AGG AGG AGC TTA TTA	<i>/</i> <i>Pse-3</i>	B2	Haley et al., 1994 Melotto et al., 1996 Fourie et al., 2004
ROC11	C11	BCMV	420 trans	CCA ATT CTC TTT CAC TTG TAA CC GCA TGT TCC AGC AAA CC	<i>bc-3</i>	B6	Johnson et al., 1997
SBD5	BD5	BCMV	1250 cis	GTG CGG AGA GGC CAT CCA TTG GTG GTG CGG AGA GTT TCA GTG TTG ACA	<i>bc-1²</i>	B3	Miklas et al., 2000a
SR2	R2	Bean Golden Yellow Mosaic Virus (BGYMV)	530 / 570 codominant	CAC AGC TGC CCT AAC AAA AT CAC AGC TGC CAC AGG TGG GA	<i>bgm-1</i>		Urrea et al., 1996 Beebe, PC, 1996
SW12	W12	BGYMV	700 cis	TGG GCA GAA GTT CTA GCA TGT GGC TGG GCA GAA GCA CAG TAT GAT TTG	Major QTL (DOR 364)	B4	Miklas et al., 2000c Singh et al. 2000
SAS8	AS08	Beet curly top virus (BCTV)	1550 cis	GGC TGC CAG TAT CTT GTC TAA CAC C GGC TGC CAG TGA CGC AAT TCT GCA G	<i>Bct</i>	B7	Larsen and Miklas, 2004
SK14	K14	Rust	620 cis	CCC GCT ACA CAC CAA TAC CTG CCC GCT ACA CTT GAT AAA ATG TTA G	<i>Ur-3</i>	B11*	Haley et al. 1994 Nemchin. & Stavely, 1998 *Miklas et al., 2002

SA14	A14	Rust	1079 / 800 codominant	CTA TCT GCC ATT ATC AAC TCA AAC GTG CTG GGA AAC ATT ACC TAT T	<i>Ur-4</i>	B6*	Miklas et al. 1993 Miene et al. 2004 *Miklas et al., 2002
SI19	I19	Rust	460 cis	AAT GCG GGA GAT ATT AAA AGG AAA G AAT GCG GGA GTT CAA TAG AAA AAC C	<i>Ur-5</i>	B4*	Haley et al., 1993 Melotto et al., 1998 *Miklas et al., 2000c
SBC6	BC06	Rust	308 cis	GAA GGC GAG AAG AAA AAG AAA AAT GAA GGC GAG AGC ACC TAG CTG AAG	<i>Ur-6</i>	B11*	Park et al., 2003b, 2004b *Miklas et al. 2002
SAD12	AD12	Rust	537 cis	AAG AGG GCG TGA GAT CGT CG AAG AGG GCG TCT TGA AGG TT	<i>Ur-7</i>	B11	Park et al., 2003a, 2004a
SAE19	AE19	Rust	890 trans	CAG TCC CTG ACA ACA TAA CAC C CAG TCC CTA AAG TAG TTT GTC CCT A	<i>Ur-11</i>	B11*	Johnson et al., 1995 Queiroz et al., 2004c *Miklas et al., 2002
UR11- GT2	GT02	Rust	450 cis	CGC ACT TAG GAG CAC AAA TGG TGG GTC CCA TAT TTT G	<i>Ur-11</i>	B11*	Boone et al., 1999 *Miklas et al., 2002
SF10	F10	Rust	1072 cis	GGA AGC TTG GTG AGC AAG GA GGA AGC TTG GCT ATGATG GT	Ouro Negro	B4*	Correa et al., 2000 *Miklas et al., 2002
SBA8	BA8	Rust	530 cis	CCA CAG CCG ACG GAG GAG GCC ATG TTT TTT GTC CCC	Ouro Negro	B4*	Correa et al., 2000 *Miklas et al., 2002

SF10	F10	Rust	1072 cis	GGA AGC TTG GTG AGC AAG GA GGA AGC TTG GCT ATGATG GT	<i>Co-10</i>	B4	Correa et al., 2000 Alzate-Marin et al., 2003
Phs	Phaseolin 'T' & 'S' alleles	White Mold & CBB	Multiple	AGC ATA TTC TAG AGG CCT CC GCT CAG TTC CTC AAT CTG TTC	Major QTL (G 122) & (BAT 93)	B7	Kami et al., 1995 Nodari et al., 1993 Miklas et al., 2001
SAU5	AU05	White mold	1350 cis	GAG CTA CCG TCA GTT TAC TAA GAG CTA CCG TGG CTT TTT TCT	QTL (minor) NY6020-4	B6	Miklas et al., 2003
SS18	S18	White mold	1650 cis	CTG GCG AAC TGT ACA TGC AAC ATA C CTG GCG AAC TGA TTC ATA CAT TTT G	QTL (major) NY6020-4	B8	Miklas et al., 2003
	SE _{ACT} /M _{CCA}	Anthracnose	codominant		<i>Co-1²</i>	B1	Vallejo and Kelly, 2002

SCAreoli	H20	Anthracnose	1000	GGG AGA CAT CCA TCA GAC AAC TCC GGG AGA CAT CTT CAT TTG ATA TGC	<i>Co-2</i>	B11	Geffroy et al., 1998 Adam-Blondon et al., 1994
SY20	Y20	Anthracnose	830 cis	AGC CGT GGA AGG TTG TCA T CCG TGG AAA CAA CAC ACA AT	<i>Co-4</i>	B8*	Queiroz et al., 2004b *Kelly et al., 2003
SC08	C08	Anthracnose	910 cis	AGA ATG CCT TTA GCT GTT GG CAG AGA GGC TAG GCT TAT CG	<i>Co-4</i>	B8*	Queiroz et al., 2004b *Kelly et al., 2003
SAS13	AS13	Anthracnose	950 cis	CAC GGA CCG AAT AAG CCA CCA ACA CAC GGA CCG AGG ATA CAG TGA AAG	<i>Co-4²</i>	B8*	Young et al., 1998 *Kelly et al., 2003
SH18	H18	Anthracnose	1100 cis	CCA GAA GGA GCT GAT AGT ACT CCA CAA C GGT AGG CAC ACT GAT GAA TCT CAT GTT GGG	<i>Co-4²</i>	B8*	Awale and Kelly, 2001 *Kelly et al., 2003
SBB14	BB14	Anthracnose	1150/1050 codominant	GTG GGA CCT GTT CAA GAA TAA TAC GTG GGA CCT GGG TAG TGT AGA AAT	<i>Co-4²</i>	B8*	Awale and Kelly, 2001 *Kelly et al., 2003

SAB3	AB-3	Anthracnose	400 cis	TGG CGC ACA CAT AAG TTC TCA CGG TGG CGC ACA CCA TCA AAA AAG GTT	<i>Co-5</i>		Vallejo and Kelly, 2001
SZ20	Z20	Anthracnose	845 cis	ACC CCT CAT GCA GGT TTT TA CAT AAT CCA TTC ATG CTC ACC	<i>Co-6</i>	B7*	Queiroz et al., 2004b *Kelly et al., 2003
SZ04	Z04	Anthracnose	567 trans	GGC TGT GCT GAT TAA TTC TGG TGC TCA TTT TAT AAT GGA GAA AAA	<i>Co-6</i>	B7*	Queiroz et al., 2004b *Kelly et al., 2003
SB12	B-12	Anthracnose	350 cis	CCT TGA CGC ACC TCC ATG TTG ACG ATGGG TTG GCC	<i>Co-9</i>	B4	Mendez de Vigo et al., 2002
SH13	H13	Angular leaf spot ALS	520 cis	GAC GCC ACA CCC ATT ATG TT GCC ACA CAG ATG GGA GCT TTA	<i>Phg-1</i>		Queiroz et al., 2004a
SN02	N02	ALS	890 cis	ACC AGG GGC ATT ATG AAC AG ACC AGG GGC AAC ATA CTA TG	<i>Phg-2</i>	B8*	Nietsche et al., 2000 *Miklas, PC, 2002

	E-ACA/M-CTT ₃₃₀	ALS	280 / 305 codominant	CTT GTT CTG AGT CAT TTA CCT TGC GAA TTC ACA GTC CAA ACT CTA ATC	G 10474 Dominant gene		Mahuku et al., 2004
SAA19	AA19	ALS	650 cis	TGA GGC GTG TCA ATG GAT ATA A GAG GCG TGT TGA TAA TTC TGG	Ouro Negro dominant gene		Queiroz et al., 2004a
SBA16	BA16	ALS	560 cis	TTC CAC GTC TAT TTT GCA TCA CAC GCA TCA CGC AGA ACT	Ouro Negro dominant gene		Queiroz et al., 2004a
SM02	M02	ALS	460 cis	CAA CGC CTC ATT AAA TTG GA CGC CTC TAA ACG GGA GAA AC	Ouro Negro dominant gene		Queiroz et al., 2004a
SW6-800R	OP W-06	Bean pod weevil, <i>Apion godmani</i>	520 cis	AGGCCCCGATGCCCCCTTAT TGGAGTCGGTCAAACCCATGTT	Agm? J-117	B11	Blair et al., 2006

SCAR Name	PCR	Protocol
SAP6	34 cycles of 10s at 94°C, 40s at 55°C, and 120s at 72°C; followed by one cycle of 5 minutes at 72°C	
BC409	34 cycles of 10s at 94°C, 60s at 70°C, and 120s at 72°C; followed by one cycle of 5 minutes at 72°C	
SU91	34 cycles of 10s at 94°C, 40s at 58°C, and 120s at 72°C; followed by one cycle of 5 minutes at 72°C	
BC420/LG5	35 cycles of 30s at 94°C, 30s at 50°C, and 60s at 72°C; followed by one cycle of 5 minutes at 72°C	
R7313	34 cycles of 10s at 94°C, 40s at 60°C, and 120s at 72°C; followed by one cycle of 5 minutes at 72°C	
R4865	34 cycles of 10s at 94°C, 40s at 60°C, and 120s at 72°C; followed by one cycle of 5 minutes at 72°C	
SB10	94°C 5 min for 1 cycle, 30 cycles of 10s at 94°C, 40s at 65°C, and 120s at 72°C; followed by one cycle of 5 minutes at 72°C	
SW13	34 cycles of 10s at 94°C, 40s at 67°C, and 120s at 72°C; followed by one cycle of 5 minutes at 72°C	
ROC11	34 cycles of 10s at 94°C, 40s at 55°C, and 120s at 72°C; followed by one cycle of 5 minutes at 72°C	
SBD5	34 cycles of 10s at 94°C, 40s at 65°C, and 120s at 72°C; followed by one cycle of 5 minutes at 72°C	
SR2	34 cycles of 10s at 94°C, 40s at 60°C, and 120s at 72°C; followed by one cycle of 5 minutes at 72°C 60°C annealing = codominant; 65°C annealing = dominant	
SW12	1 cycle of 60s at 94°C; 30 cycles of 30s at 94°C, 30s at 70°C, and 60s at 72°C; followed by one cycle of 5 minutes at 72°C	
SAS8	1 cycle of 5 min at 94°C; 30 cycles of 60s at 94°C, 60s at 68°C, and 60s at 72°C; followed by one cycle of 7 minutes at 72°C	
SK14	34 cycles of 10s at 94°C, 40s at 63°C, and 120s at 72°C; followed by one cycle of 5 minutes at 72°C	
SA14	1 cycle of 5 min at 94°C; 35 cycles of 60s at 94°C, 60s at 55°C, and 90s at 72°C; followed by one cycle of 5 minutes at 72°C	
SI19	34 cycles of 10s at 94°C, 40s at 67°C, and 120s at 72°C; followed by one cycle of 5 minutes at 72°C	
SBC6	94°C 2 min; 30 cycles of 30s at 94°C, 60s at 59 or 65°C?, and 120s at 72°C; followed by one cycle of 7 minutes at 72°C	
SAD12	Annealing temperature of 71°C	
SAE19	94°C 5 min; 35 cycles of 15s at 94°C, 60s at 58°C, and 90s at 72°C; followed by one cycle of 7 minutes at 72°C	

UR11-GT2	60°C annealing = codominant; 65°C annealing = dominant
SF10	1 cycle of 94°C of 3 min: 35 Cycles at 15s at 94°C, 60s at 65°C and 90s at 72°C; followed by 1 cycle of 7 min at 72°C
SBA8	1 cycle of 94°C of 3 min; 35 Cycles at 15s at 94°C, 60s at 65°C and 90s at 72°C; followed by 1 cycle of 7 min at 72°C
Phs	34 cycles of 10s at 94°C, 40s at 50°C, and 120s at 72°C; followed by one cycle of 5 minutes at 72°C
SAU5	34 cycles of 10s at 94°C, 40s at 60°C, and 120s at 72°C; followed by one cycle of 5 minutes at 72°C
SS18	34 cycles of 10s at 94°C, 40s at 63°C, and 120s at 72°C; followed by one cycle of 5 minutes at 72°C
SCAreoli	58°C annealing temperature followed by <i>DraI</i> digestion
SY20	35 cycles of 30s at 94°C, 60s at 65°C, and 90s at 72°C
SC08	35 cycles of 30s at 94°C, 60s at 65°C, and 90s at 72°C
SAS13	34 cycles of 10s at 94°C, 144s at 72°C; followed by one cycle of 5 minutes at 72°C
SH18	34 cycles of 10s at 94°C, 40s at 62°C, and 120s at 72°C; followed by one cycle of 5 minutes at 72°C
SBB14	34 cycles of 10s at 94°C, 40s at 67°C, and 120s at 72°C: followed by one cycle of 5 minutes at 72°C
SAB3	1 cycle of 3 minutes at 94°C; 30 cycles of 10s at 94°C, 30s at 65°C, 2 minutes at 72°C; followed by 1 cycle of 5 minutes at 72°C
SZ20	35 cycles of 30s at 94°C, 60s at 60°C, and 90s at 72°C
SZ04	45 cycles of 30s at 94°C, 120s at 45°C, and 90s at 72°C
SB12	1 cycle of 94°C of 2 min: 35 Cycles of 60s at 94°C, 60s at 68°C, and 60s at 72°C; followed by 1 cycle of 7 min at 72°C
SF10	1 cycle of 94°C of 3 min: 35 Cycles at 15s at 94°C, 60s at 65°C and 90s at 72°C; followed by 1 cycle of 7 min at 72°C
SH13	35 cycles of 30s at 94°C, 60s at 59°C, and 90s at 72°C
SN02	30 Cycles of 30s at 94°C, 60s at 65°C, and 90s at 72°C
G10474	1 cycle of 94°C of 5 min: 35 Cycles of 30s at 94°C, 45s at 60°C, and 30s at 72°C; followed by 1 cycle of 10 min at 72°C
SAA19	35 cycles of 30s at 94°C, 60s at 56°C, and 90s at 72°C
SBA16	35 cycles of 30s at 94°C, 60s at 58°C, and 90s at 72°C
SM02	35 cycles of 30s at 94°C, 60s at 58°C, and 90s at 72°C

References for SCAR list:

- Adam-Blondon, A.F., M. Seignac, H. Bannerot, and M. Dron. 1994. SCAR, RAPD, and RFLP markers linked to the dominant gene (Are) conferring resistance to anthracnose. *Theor. Appl. Genet.* 88:865-870.
- Alzate-Marin, A.L., M.R. Costa, K.M. Arruda, E.G. Barros, and M.A. Moreira. 2003. Characterization of the anthracnose resistance gene present in Ouro Negro (Honduras 35) common bean cultivar. *Euphytica* 133:165-169.
- Ariyaratne, H.M., D.P. Coyne, G. Jung, P.W. Skroch, A.K. Vidaver, J.R. Steadman, P.N. Miklas, and M.J. Bassett. 1999. Molecular mapping of disease resistance genes for halo blight, common bacterial blight, and bean common mosaic virus in a segregating population of common bean. *J. Am. Soc. Hort. Sci.* 124:654-662.
- Awale, H.E., and J.D. Kelly. 2001. Development of SCAR markers linked to *Co-4* gene in common bean. *Annu Rept. Bean Improv. Coop.* 44:119-120.
- Bai, Y., T.E. Michaels, and K.P. Pauls. 1997. Identification of RAPD markers linked to common bacterial blight resistance genes in *Phaseolus vulgaris* L. *Genome* 40:544-551.
- Beattie, A., T.E. Michaels, and K.P. Paul. 1998. An efficient reliable method to screen for common bacterial blight (CBB) resistance in *Phaseolus vulgaris* L. *Annu. Rept. Bean Improv. Coop.* 41:53-54.
- Blair, M.W., C. Muñoz, R. Garza and C. Cardona. 2006. Molecular mapping of genes for resistance to the bean pod weevil (*Apion godmani* Wagner) in common bean. *Theor. Appl. Genet.* 112:913-923.
- Boone, W.E., J.R. Stavely, and N.F. Weeden. 1999. Development of a sequence-tagged site (STS) marker for *Ur-11*, a gene conferring resistance to the bean rust fungus, *Uromyces appendiculatus*. *Annu. Rept. Bean Improv. Coop.* 42:33-34.
- Corrêa, R.X., M.R. Costa, P.I. Good-God, V.A. Ragagnin, F.G. Faleiro, M.A. Moreira, and E.G. de Barros. 2000. Sequence characterized amplified regions linked to rust resistance genes in the common bean. *Crop Sci.* 40:804-807.
- Fourie, D., P.N. Miklas, and H.M. Ariyaratne. 2004. Genes conditioning halo blight resistance to races 1, 7, and 9 occur in a tight cluster. *Annu. Rep. Bean Improv. Coop.* 47:103-104.

- Geffroy, V., F. Creusot, J. Falquet, M. Seignac, A.F. Adam-Blondon, H. Bannerot, P. Gepts, and M. Dron. 1998. A family of LRR sequences at the *Co-2* locus for anthracnose resistance in *Phaseolus vulgaris* and its potential use in marker-assisted selection. *Theor. Appl. Genet.* 96:494-502.
- Haley, S.D., L. Afanador, and J.D. Kelly. 1994. Identification and application of a random amplified polymorphic DNA marker for the */* gene (potyvirus resistance) in common bean. *Phytopathology* 84:157-160.
- Haley, S.D., L.K. Afanador, P.N. Miklas, J.R. Stavely, and J.D. Kelly. 1994. Heterogeneous inbred populations are useful as sources of near-isogenic lines for RAPD marker localization. *Theor. Appl. Genet.* 88: 337-342.
- Haley, S.D., P.N. Miklas, J.R. Stavely, J. Byrum, and J.D. Kelly. 1993. Identification of RAPD markers linked to a major rust resistance gene block in common bean. *Theor. Appl. Genet.* 86:505-512.
- Johnson, W.C., P. Guzman, D. Mandala, A.B.C. Mkandawire, S. Temple, R.L. Gilbertson, and P. Gepts. 1997. Molecular tagging of the *bc-3* gene for introgression into Andean common bean. *Crop Sci.* 37:248-254.
- Johnson, E., P.N. Miklas, J.R. Stavely, and J.C. Martinez-Cruzado. 1995. Coupling- and repulsion-phase RAPDs for marker-assisted selection of PI 181996 rust resistance in common bean. *Theor. Appl. Genet.* 90:659-664.
- Jung, G., Skroch, P.W., Nienhuis, J., Coyne, D.P., Arnaud-Santana, E., Ariyaratne, H.M., Marita, J.M. 1999. Confirmation of QTL associated with common bacterial blight resistance in four different genetic backgrounds in common bean. *Crop Sci.* 39, 1448-1455.
- Kami, J., V.B. Velasquez, D.G. Debouck, and P. Gepts. 1995. Identification of presumed ancestral DNA sequences of phaseolin in *Phaseolus vulgaris*. *Proc. Natl. Acad. Sci.* 92:1101-1104.
- Kelly, J.D., P. Gepts, P.N. Miklas, and D.P. Coyne. 2003. Tagging and mapping of genes and QTL and molecular marker-assisted selection for traits of economic importance in bean and cowpea. *Field Crops Research* 82:135-154.
- Kelly, J.D. and P.N. Miklas. 1999. Marker-assisted selection. p.93-123. *In* S. Singh (ed.) *Common bean improvement in the twenty-first century*. Kluwer Academic Publ., The Netherlands.
- Larsen, R.C., and P.N. Miklas. 2004. Generation and molecular mapping of a SCAR marker linked with the *Bct* gene for resistance to *beet curly top virus* in common bean. *Phytopathology* 94:320-325.
- Mahuku, G., C. Montoya, M.A. Henríquez, C. Jara, H. Teran, and S. Beebe. 2004. Inheritance and characterization of angular leaf spot resistance gene present in common bean accession G 10474 and identification of an AFLP marker linked to the resistance gene. *Crop Sci.* 44:1817-1824.
- Meine, C.M.S., R Naidoo, and M.M. Liebenberg. 2004. Conversion of the RAPD marker for *Ur-4* to a co-dominant SCAR marker SA14_{1079/800}. *Annu. Rep. Bean Improv. Coop.* 47:261-262.

- Melotto, M., L. Afanador, and J.D. Kelly. 1996. Development of a SCAR marker linked to the */* gene in common bean. *Genome* 39:1216-1219.
- Melotto, M. and J.D. Kelly. 1998. SCAR markers linked to major disease resistance genes in common bean. *Annu. Rept. Bean Improv. Coop.* 41:64-65.
- Mendez de Vigo, B., C. Rodriguez, A. Paneda, R. Giraldez, and J.J. Ferreira. 2002. Development of a SCAR marker linked to *Co-9* in common bean. *Annu. Rep. Bean Improv. Coop.* 45:116-117.
- Miklas, P.N., R. Delorme, and R.H. Riley. 2003. Identification of QTL conditioning resistance to white mold in a snap bean population. *J. Am. Soc. Hort. Sci.* 128:564-570.
- Miklas, P.N., E. Johnson, V. Stone, J.S. Beaver, C. Montoya, and M. Zapata. 1996. Selective mapping of QTL conditioning disease resistance in common bean. *Crop Sci.* 36:1344-1351.
- Miklas, P.N., W.C. Johnson, R. Delorme, R.H. Riley, and P. Gepts. 2001. Inheritance and QTL analysis of physiological resistance to white mold in common bean G122. *Crop Sci.* 41:309-315.
- Miklas, P.N., R. Larsen, K. Victry, R. Delorme, C. Marma, R.H. Riley, & James D. Kelly. 2000a. Marker-assisted selection for the *bc-1²* gene for resistance to BCMV and BCMNV in common bean. *Euphytica* 116:211-219.
- Miklas, P.N., M.A. Pastor-Corrales, G. Jung, D.P. Coyne, J.D. Kelly, P.E. McClean, and P. Gepts. 2002. Comprehensive linkage map of bean rust resistance genes. *Annu. Rep. Bean Improv. Coop.* 45:125-129.
- Miklas, P.N., J.R. Smith, R. Riley, K.F. Grafton, S.P. Singh, G. Jung, and D. P. Coyne. 2000b. Marker-assisted breeding for pyramided resistance to common bacterial blight in common bean. *Annu. Rep. Bean Improv. Coop.* 43:39-40.
- Miklas, P.N., J.R. Stavely, and J.D. Kelly. 1993. Identification and potential use of a molecular marker for rust resistance in common bean. *Theor. Appl. Genet.* 85:745-749.
- Miklas, P.N., V. Stone, M.J. Daly, J.R. Stavely, J.R. Steadman, M.J. Bassett, R. Delorme, and J.S. Beaver. 2000c. Bacterial, fungal, and viral disease resistance loci mapped in a recombinant inbred common bean population ('Dorado'/XAN 176). *J. Am. Soc. Hort. Sci.* 125: 476-481.
- Nemchinova, Y.P., and J.R. Stavely. 1998. Development of SCAR primers for the *Ur-3* rust resistance gene in common bean. *Phytopathology* 88:S67.
- Nietsche, S., A. Borem, G.A. Carvalho, R.C. Rocha, T.J. Paula, E.G. de Barros, M.A. Moreira. 2000. RAPD and SCAR markers linked to a gene conferring resistance to angular leaf spot in common bean. *J. of Phytopathology* 148:117-121.

- Nodari, R.O., S.M. Tsai, P. Guzman, R.L. Gilbertson, and P. Gepts. 1993. Towards an integrated linkage map of common bean. III. Mapping genetic factors controlling host-bacteria interactions. *Genetics* 134:341-350.
- Park, S.O., D.P. Coyne, J.R. Steadman. 2004a. Development of a SCAR marker linked to the *Ur-7* gene in common bean. *Annu. Rep. Bean Improv. Coop.* 47:269-270.
- Park, S.O., D.P. Coyne, J.R. Steadman, K.M. Crosby, and M.A. Brick. 2004b. RAPD and SCAR markers linked to the *Ur-6* Andean gene controlling specific rust resistance in common bean. *Crop Sci.* 44:1799-1807.
- Park, S.O., D.P. Coyne, J.R. Steadman, and P.W. Skroch. 2003a. Mapping of the *Ur-7* gene for specific resistance to rust in common bean. *Crop Sci.* 43:1470-1476.
- Park, S.O., K.M. Crosby, D.P. Coyne, and J.R. Steadman. 2003b. Development of a SCAR marker linked to the *Ur-6* gene for specific rust resistance in common bean. *Annu Rep. Bean Improv. Coop.* 46:189-190.
- Pedraza, F., G. Gallego, S. Beebe, and J. Tohme. 1997. Marcadores SCAR y RAPD para la resistencia a la bacteriosis comun (CBB). p.130-134. En Singh, S.P. y O. Voysest (eds.). *Taller de mejoramiento de frijol para el Siglo XXI: Bases para una estrategia para America Latina*. 559 pp. CIAT, Cali, Colombia.
- Queiroz, V.T., C.S. Sousa, M.R. Costa, D.A. Sanglad, K.M.A. Arruda, T.L.P.O. Souza, V.A. Ragagnin, E.G. Barros, and M.A. Moreira. 2004a. Development of SCAR markers linked to common bean angular leaf spot resistance genes. *Annu. Rep. Bean Improv. Coop.* 47:237-238.
- Queiroz, V.T., C.S. Sousa, M.R. Costa, D.A. Sanglad, K.M.A. Arruda, T.L.P.O. Souza, V.A. Ragagnin, E.G. Barros, and M.A. Moreira. 2004b. Development of SCAR markers linked to common bean anthracnose resistance genes *Co-4* and *Co-6*. *Annu. Rep. Bean Improv. Coop.* 47:249-250.
- Queiroz, V.T., C.S. Sousa, T.L.P.O. Souza, M.R. Costa, D.A. Sanglad, V.A. Ragagnin, E.G. Barros, and M.A. Moreira. 2004c. SCAR marker linked to the common bean rust resistance gene *Ur-11*. *Annu. Rep. Bean Improv. Coop.* 47:271-272.
- Singh, S.P., F.J. Morales, P.N. Miklas, and H. Terán. 2000. Selection for bean golden mosaic resistance in intra- and inter-racial bean populations. *Crop Sci.* 40:1565-1572.
- Urrea, C.A., P.N. Miklas, J.S. Beaver, and R.H. Riley. 1996. A codominant randomly amplified polymorphic DNA (RAPD) marker useful for indirect selection of BGMV resistance in common bean. *J. Amer. Soc. Hort. Sci.* 121:1035-1039.
- Vallejo, V., and J.D. Kelly. 2001. Development of a SCAR marker linked to *Co-5* gene in common bean. *Annu Rep. Bean Improv. Coop.* 44:121-122.
- Vallejo, V., and J.D. Kelly. 2002. The use of AFLP analysis to tag the *Co12* gene conditioning resistance to bean anthracnose. *Plant Animal Genome X Conference 2002*, San Diego, CA.

Young, R.A. and J.D. Kelly. 1997. RAPD markers linked to three major anthracnose resistance genes in common beans. *Crop Sci.* 37:940-946

Young, R.A., M. Melotto, R.O. Nodari, and J.D. Kelly. 1998. Marker assisted dissection of the oligogenic anthracnose resistance in common bean cultivar, G 2333. *Theor. Appl. Genet.* 96:87-94.

Yu, K., S.J. Park, V. Poysa. 2000. Marker-assisted selection of common beans for resistance to common bacterial blight: efficiency and economics. *Plant Breed.* 119:411-416.